

Front Side

150 mm

25 mm

25 mm

200 mm



KETOCONAZOLE- ketoconazole cream


KETOCONAZOLE CREAM

Rx only

DESCRIPTION

Ketoconazole Cream, 2% contains the broad-spectrum synthetic antifungal agent, ketoconazole 2%. Each gram, for topical administration, contains ketoconazole 20 mg and is formulated in an aqueous cream vehicle consisting of propylene glycol, purified water, cetyl alcohol, stearyl alcohol, isopropyl myristate, sorbitan monostearate, polysorbate 60, polysorbate 80, and sodium sulfite, anhydrous:

Ketoconazole is *cis*-1-acetyl-4-[4-[[2-(2,4-dichlorophenyl)-2-(1*H*-imidazol-1-ylmethyl)-1,3-dioxolan-4-yl]methoxy]phenyl]piperazine and has the following structural formula:



C₂₆H₂₈Cl₂N₄O₄ M.W. 531.44

CLINICAL PHARMACOLOGY

When Ketoconazole Cream, 2% was applied dermally to intact or abraded skin of Beagle dogs for 28 consecutive days at a dose of 80 mg, there were no detectable plasma levels using an assay method having a lower detection limit of 2 ng/ml.

After a single topical application to the chest, back and arms of normal volunteers, systemic absorption of ketoconazole was not detected at the 5 ng/ml level in blood over a 72-hour period.

Two dermal irritancy studies, a human sensitization test, a phototoxicity study and a photoallergy study conducted in 38 male and 62 female volunteers showed no contact sensitization of the delayed hypersensitivity type, no irritation, no phototoxicity and no photoallergic potential due to Ketoconazole Cream, 2%.

Microbiology

Ketoconazole is a broad spectrum synthetic antifungal

agent which inhibits the *in vitro* growth of the following common dermatophytes and yeasts by altering the permeability of the cell membrane: dermatophytes: *Trichophyton rubrum*, *T. mentagrophytes*, *T. tonsurans*, *Microsporium canis*, *M. audouini*, *M. gypseum* and *Epidermophyton floccosum*; yeasts: *Candida albicans*, *Malassezia ovale* (*Pityrosporum ovale*) and *C. tropicalis*; and the organism responsible for tinea versicolor, *Malassezia furfur* (*Pityrosporum orbiculare*). Only those organisms listed in the INDICATIONS AND USAGE Section have been proven to be clinically affected. Development of resistance to ketoconazole has not been reported.

Mode of Action

In vitro studies suggest that ketoconazole impairs the synthesis of ergosterol, which is a vital component of fungal cell membranes. It is postulated that the therapeutic effect of ketoconazole in seborrheic dermatitis is due the reduction of *M. ovale*, but: this has not been proven.

INDICATIONS AND USAGE

Ketoconazole Cream, 2% is indicated for the topical treatment of tinea corporis, tinea cruris and tinea pedis caused by *Trichophyton rubrum*, *T. mentagrophytes* and *Epidermophyton floccosum*; in the treatment of tinea (pityriasis) versicolor caused by *Malassezia furfur* (*Pityrosporum orbiculare*); and in the treatment of cutaneous candidiasis caused by *Candida spp.* and in the treatment of seborrheic dermatitis.

CONTRAINDICATIONS

Ketoconazole Cream, 2% is contraindicated in persons who have shown hypersensitivity to the active or excipient ingredients of this formulation.

WARNINGS

Ketoconazole Cream, 2% is not for ophthalmic use.

Ketoconazole Cream, 2% contains sodium sulfite anhydrous, a sulfite that may cause allergic-type reactions including anaphylactic symptoms and life-threatening or less severe asthmatic episodes in certain susceptible people. The overall prevalence of sulfite sensitivity in the general population is unknown and probably low. Sulfite sensitivity is seen more frequently in asthmatic than in nonasthmatic people.

PRECAUTIONS

General

If a reaction suggesting sensitivity or chemical irritation should occur, use of the medication should be discontinued. Hepatitis (1:10,000 reported incidence) and,



Back Side

150 mm

at high doses, lowered testosterone and ACTH induced corticosteroid serum levels have been seen with orally administered ketoconazole; these effects have not been seen with topical ketoconazole.

Carcinogenesis, Mutagenesis, Impairment of Fertility

A long-term feeding study in Swiss Albino mice and in Wistar rats showed no evidence of oncogenic activity. The dominant lethal mutation test in male and female mice revealed that single oral doses of ketoconazole as high as 80 mg/kg produced no mutation in any stage of germ cell development. The Ames' *Salmonella* microsomal activator assay was also negative.

Pregnancy

Teratogenic effects

Ketoconazole has been shown to be teratogenic (syndactylia and oligodactylia) in the rat when given orally in the diet at 80 mg/kg/day, (10 times the maximum recommended human oral dose). However, these effects may be related to maternal toxicity, which was seen at this and higher dose levels.

There are no adequate and well-controlled studies in pregnant women. Ketoconazole should be used during pregnancy only if the potential benefit justifies the potential risk to the fetus.

Nursing Mothers

It is not known whether Ketoconazole Cream, 2% administered topically could result in sufficient systemic absorption to produce detectable quantities in breast milk. Nevertheless, a decision should be made whether to discontinue nursing or discontinue the drug, taking into account the importance of the drug to the mother.

Pediatric Use

Safety and effectiveness in children have not been established.

ADVERSE REACTIONS

During clinical trials 45 (5%) of 905 patients treated with Ketoconazole Cream, 2% and 5 (2.4%) of 208 patients treated with placebo reported side effects consisting mainly of severe irritation, pruritus and stinging. One of the patients treated with ketoconazole cream developed a painful allergic reaction.

In worldwide postmarketing experience, rare reports of contact dermatitis have been associated with ketoconazole cream or one of its excipients, namely sodium sulfite or propylene glycol.

To report SUSPECTED ADVERSE REACTIONS, contact Encube Ethicals Private Limited at 1-833-285-4151 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

DOSAGE AND ADMINISTRATION

Cutaneous candidiasis, tinea corporis, tinea cruris, tinea pedis, and tinea (pityriasis) versicolor: It is recommended

that Ketoconazole Cream, 2% be applied once daily to cover the affected and immediate surrounding area. Clinical improvement may be seen fairly soon after treatment is begun; however, candidal infections and tinea cruris and corporis should be treated for two weeks in order to reduce the possibility of recurrence. Patients with tinea versicolor usually require two weeks of treatment. Patients with tinea pedis require six weeks of treatment.

If a patient shows no clinical improvement after the treatment period, the diagnosis should be redetermined.

Seborrheic dermatitis: Ketoconazole cream, 2% should be applied to the affected area twice daily for four weeks or until no clinical clearing. If a patient shows no clinical improvement after the treatment period, the diagnosis should be redetermined.

HOW SUPPLIED

Ketoconazole Cream, 2% is supplied in 15, 30, and 60 gram tubes.

15 g tube (21922-025-04)

30 g tube (21922-025-05)

60 g tube (21922-025-07)

STORAGE

Store at 20° to 25°C (68° to 77°F) [See USP Controlled Room Temperature].

Mfg. Lic. No. 361

Manufactured by:

Encube Ethicals Pvt. Ltd.

Plot No. C-1, Madkaim Industrial Estate,

Madkaim, Post: Mardol, Ponda,

Goa - 403 404, India.

Distributed by:

Encube Ethicals, Inc.

200 Meredith Avenue, Suite 101A

Durham, NC 27713 USA

Rev.: 03


August 2020



EKC5/00

Revision

Rev. No.	Date	Artist	Correction
2	29.07.2020	Pranita Graphics	1) Added (.) after Lic word in Mfg. Lic. No. 2) Revised Rev No. 02 to 03 & March 2020 to August 2020.
3	03.08.2020	Pranita Graphics	1) Artwork code Revised EKC3C/00 to EKC5C/00
4	18.08.2020	Pranita Graphics	1) Changed folding Size 20 mm to 25 mm 2) Change in text positioning, alignment due to Chang in folding size
5	20.08.2020	Pranita Graphics	1) Added full stop after word <i>spp.</i> in INDICATIONS AND USAGE Para.
6	06.04.2021	Pranita Graphics	1) Changed size 120 x 200 mm to 150 x 200 mm as per Production Department Req. 2) Kept space before Dosage and Administration para 3) Added space in between "August 2020" 4) corrected Artwork code EKC5C/00 to EKC5/00

 Encube	Product	Ketoconazole Cream, 2% Pack Insert			Colours Used
	Artwork Code	EKC5/00		<input checked="" type="checkbox"/>	Black
	SAP Code	206759			
	Pharma code	22			
	Size	150 x 200 mm Folding Size: 25 mm			
	Material	Maplitho			
	Grammage	-			
	Varnish	-		<input type="checkbox"/>	Keyline (not to print)
	Spl Req.	NA			
	Changes history after submission		Size change		Minimum Font Size: 7.5 pt
				Maximum Font Size: 11.3pt	
				NV/08.04.2021	
Prepared by:		Reviewed by:			Approved by:
Pkg. Devp. Dept.	Production Dept.	Engineering Dept.	RA (HO)	QA/RA (GOA)	Head Pkg. Devp.